



● RESOURCE

Acute health risks to community hand-pumped groundwater supplies following Cyclone Idai flooding

Author(s)

Rivett, Michael O. Tremblay Lévesque, Laurent-Charles Carter, Ruth Thetard, Rudi C.H. Tengtenga, Morris Phoya, Ann Mbalame, Emma Mchilikizo, Edwin Kumwenda, Steven Mleta, Prince Addison, Marc J. Kalin, Robert M.

Description / Abstract

This longitudinal flood-relief study assessed the impact of the March 2019 Cyclone Idai flood event on *E. coli* contamination of hand-pumped boreholes in Mulanje District, Malawi. It established the microbiological water-quality safety of 279 community supplies over three phases, each comprising water-quality survey, rehabilitation and treatment verification monitoring. Phase 1 contamination three months after Idai was moderate, but likely underestimated. Increased contamination in Phase 2 at 9 months and even greater in Phase 3, a year after Idai was surprising and concerning, with 40% of supplies then registering *E. coli* contamination and 20% of supplies deemed 'unsafe'. Without donor support for follow-up interventions, this would have been missed by a typical single-phase flood-relief activity. Contamination rebound at boreholes successfully treated months earlier signifies a systemic problem from persistent sources intensified by groundwater levels likely at a decade high. Problem extent in normal, or drier years is unknown due to absence of routine monitoring of water point *E. coli* in Malawi. Statistical analysis was not conclusive, but was indicative of damaged borehole infrastructure and increased near-borehole pit-latrines being influential. Spatial analysis including groundwater flow-field definition (an overlooked sector opportunity) revealed 'hit-and-miss' contamination of safe and unsafe boreholes in proximity. Hydrogeological control was shown by increased contamination near flood-affected area and in more recent recharge groundwater otherwise of good quality. Pit latrines are presented as credible *e-coli* sources in a conceptual model accounting for heterogeneous borehole contamination, wet season influence and rebound behavior. Critical to establish are groundwater level - flow direction, hand-pump plume draw, multiple footprint latrine sources - 'skinny' plumes, borehole short-circuiting and fast natural pathway (e.g. fracture flow) and other source influences. Concerted WASH (Water, Sanitation and Hygiene) sector investment in research and policy driving national water point based *E. coli* monitoring programs are advocated.

Publication year

2021

Country

Malawi

Region

Africa

Publisher

Science of the Total Environment

Keywords

Cyclone Idai E.Coli contamination Pit latrine Water, Sanitation and Hygiene (WASH) Groundwater Management Mulanje District

Thematic Tagging

Climate Ecosystems/Nature-based solutions Gender Private Sector Transboundary Urban Water services Youth

Language English

[View resource](#)

Related IWRM Tools



Tool

Geographic Information System

C2.01

Source URL: <https://iwrmactionhub.org/resource/acute-health-risks-community-hand-pumped-groundwater-supplies-following-cyclone-idai>